

# A CORRESPONDENCE BETWEEN HIGHER ADAMS DIFFERENTIALS AND HIGHER ALGEBRAIC NOVIKOV DIFFERENTIALS AT ODD PRIMES

XIANGJUN WANG AND YU ZHANG\*

ABSTRACT. This paper studies the higher differentials of the classical Adams spectral sequence at odd primes. In particular, we follow the “cofiber of  $\tau$  philosophy” of Gheorghe, Isaksen, Wang, and Xu [Acta Math. 226 (2021), pp. 319–407; Publ. Math. Inst. Hautes Etudes Sci. 137 (2023), pp. 107–243] to show that higher Adams differentials agree with their corresponding higher algebraic Novikov differentials in a certain range.

## 1. INTRODUCTION

The computation of the stable homotopy groups of the sphere  $\pi_*(S^0)$  is one of the most important problems in homotopy theory. In recent years, a major breakthrough in this area is the work of Isaksen-Wang-Xu [8], which successfully extended the computation of the 2-primary components of  $\pi_*(S^0)$  to dimension 90. Their computation is based on the “cofiber of  $\tau$  philosophy” developed in Gheorghe-Wang-Xu [5]. The key insight of [5, 8] is that one can compute difficult higher Adams differentials from higher algebraic Novikov differentials using motivic homotopy theory. Although the method developed in [5] holds for arbitrary primes, it has only been extensively exploited for 2-primary computations. In this paper, we apply the “cofiber of  $\tau$  philosophy” to odd-primary computations.

**Some classical spectral sequences.** Throughout this paper, we let  $p$  denote a fixed odd prime. The Adams spectral sequence (ASS) and the Adams-Novikov spectral sequence (ANSS) are two of the most powerful tools to compute the  $p$ -primary component of  $\pi_*(S^0)$ .

The  $E_2$ -page of the ASS is  $Ext_{\mathcal{A}_*}^{*,*}(\mathbb{F}_p, \mathbb{F}_p)$ , where  $\mathcal{A}_*$  is the dual mod  $p$  Steenrod algebra. We can write  $\mathcal{A}_* = \mathbb{F}_p[t_1, t_2, \dots] \otimes E[\tau_0, \tau_1, \tau_2, \dots]$ , where  $\mathbb{F}_p[t_1, t_2, \dots]$  is a polynomial algebra with coefficients in  $\mathbb{F}_p$ , and  $E[\tau_0, \tau_1, \tau_2, \dots]$  is an exterior algebra with coefficients in  $\mathbb{F}_p$ . The differentials of the ASS have the form  $d_r^{Adams} : E_r^{s,t}(S) \rightarrow E_r^{s+r,t+r-1}(S)$ ,  $r \geq 2$ .

The  $E_2$ -page of the ANSS is  $Ext_{BP_*BP}^{*,*}(BP_*, BP_*)$ , where  $BP$  denotes the Brown-Peterson spectrum. We have  $BP_* := \pi_*(BP) = \mathbb{Z}_{(p)}[v_1, v_2, \dots]$ , and that  $BP_*BP = BP_*[t_1, t_2, \dots]$ , where  $\mathbb{Z}_{(p)}$  denotes the integers localized at  $p$ . The inner degrees of the generators are  $|v_n| = |t_n| = 2(p^n - 1)$ .

---

2020 *Mathematics Subject Classification.* 14F42, 55Q45, 55T15, 55T25.

Keywords: Motivic homotopy theory, Stable homotopy of spheres, Adams spectral sequences, algebraic Novikov spectral sequences.

\* Corresponding author.

The algebraic Novikov spectral sequence (algNSS) [9, 13] converges to the Adams-Novikov  $E_2$ -page. The  $E_2$ -page of the algNSS is  $Ext_{P_*}^{s,t}(\mathbb{F}_p, I^k/I^{k+1})$ , where  $I$  denotes the ideal  $(p, v_1, v_2, \dots) \subset BP_*$ , and  $P_* = BP_*BP/I = \mathbb{F}_p[t_1, t_2, \dots]$  is the  $\mathbb{F}_p$ -coefficient polynomial algebra. The differentials have the form  $d_r^{alg} : \bar{E}_r^{s,t,k} \rightarrow \bar{E}_r^{s+1,t,k+r-1}$ ,  $r \geq 2$ .

*Remark 1.1.* Here, we have re-indexed the pages of the algNSS to align with the notations in [5, 8].

The  $E_2$ -page of the Adams spectral sequence can also be computed via a spectral sequence, called the Cartan-Eilenberg spectral sequence (CESS) [2, 15]. For odd primes  $p$ , the  $E_2$ -page of the CESS coincides with the  $E_2$ -page of the algNSS up to degree shifting [9].

These four spectral sequences fit into the following square [10].

$$(1.1) \quad \begin{array}{ccc} Ext_{P_*}^{s,t}(\mathbb{F}_p, I^k/I^{k+1}) & \xrightarrow{CESS} & Ext_{\mathcal{A}_*}^{s+k,t+k}(\mathbb{F}_p, \mathbb{F}_p) \\ \downarrow algNSS & & \downarrow ASS \\ Ext_{BP_*BP}^{s,t}(BP_*, BP_*) & \xrightarrow{ANSS} & \pi_{t-s}(\widehat{S}^0) \end{array}$$

The Adams differentials  $d_r^{Adams}$  and the Adams-Novikov differentials  $d_r^{AN}$  are difficult to compute in general. Such computations might require an understanding of complicated geometric behaviors. On the other hand, the algebraic Novikov differentials  $d_r^{alg}$  are much easier to compute. This is because the entire construction of the algNSS is purely algebraic.

It is long been observed that  $d_r^{alg}$  and  $d_r^{Adams}$  are closely related. It is desirable if we could determine  $d_r^{Adams}$  differentials based on  $d_r^{alg}$  computations. The following result realizes the idea for  $r = 2$ , while the problem for general  $r > 2$  remains open.

**Theorem 1.2** (Novikov [13], Andrews-Miller [1, 10]). *Suppose  $z$  is an element in  $Ext_{\mathcal{A}_*}^{s+k,t+k}(\mathbb{F}_p, \mathbb{F}_p)$  which is detected in the CESS by  $x \in Ext_{P_*}^{s,t}(\mathbb{F}_p, I^k/I^{k+1})$ . Then,  $d_2^{Adams}(z)$  is detected by  $d_2^{alg}(x) \in Ext_{P_*}^{s+1,t}(\mathbb{F}_p, I^{k+1}/I^{k+2})$ .*

*Remark 1.3.* Indeed, one can use this approach to find new nontrivial secondary Adams differentials. See [19] for some practical examples.

**New developments using motivic homotopy theory.** For general  $r \geq 2$ , Gheorghe-Wang-Xu [5] developed the ‘‘cofiber of  $\tau$  philosophy’’ to compare  $d_r^{alg}$  with  $d_r^{Adams}$ . One key feature of their approach is the usage of motivic homotopy theory, which originated from the work of Morel and Voevodsky [12]. Motivic homotopy theory behaves very similarly to the classical homotopy theory. For example, there are motivic dual Steenrod algebras [16, 17] and motivic Adams spectral sequences (mASS) [4, 6].

The computational technique of [5] can be illustrated in the following diagram.

$$(1.2) \quad \begin{array}{ccccc} Ext_{\mathcal{A}_{*,*}^c}^{*,*,*}(\mathbb{F}_p[\tau], \mathbb{F}_p) & \longleftarrow & Ext_{\mathcal{A}_{*,*}^c}^{*,*,*}(\mathbb{F}_p[\tau], \mathbb{F}_p[\tau]) & \longrightarrow & Ext_{\mathcal{A}_*}^{*,*}(\mathbb{F}_p, \mathbb{F}_p) \\ \downarrow mASS & & \downarrow mASS & & \downarrow ASS \\ \pi_{*,*}(\widehat{S}^{0,0}/\tau) & \longleftarrow & \pi_{*,*}(\widehat{S}^{0,0}) & \longrightarrow & \pi_*(\widehat{S}^0) \end{array}$$

They considered a map  $\tau : \widehat{S^{0,-1}} \rightarrow \widehat{S^{0,0}}$  between  $p$ -completed motivic sphere spectra. We let  $\widehat{S^{0,0}}/\tau$  denote the cofiber of the map  $\tau$ . In diagram (1.2), the left column is the mASS for  $\widehat{S^{0,0}}/\tau$ , the middle column is the mASS for  $\widehat{S^{0,0}}$ , and the right column is the classical ASS. The top horizontal maps are maps of spectral sequences. Gheorghe-Wang-Xu [5] proved the left column is isomorphic to the algNSS for any prime  $p$ . Hence diagram (1.2) provides a zig-zag diagram to compare higher Adams differentials with their corresponding higher algebraic Novikov differentials.

For  $p = 2$ , Isaksen-Wang-Xu [8] used the strategy above to determine  $d_r^{Adams}$  based on computer generated  $d_r^{alg}$  data. This enabled them to successfully extend the computation of 2-primary stable homotopy groups from around 60 stems to around 90 stems.

**Our main results.** In this paper, we follow the ‘‘cofiber of  $\tau$  philosophy’’ but focus on the case when  $p$  is an odd prime. For odd prime  $p$ , the motivic Steenrod algebra has a simpler form and the diagram (1.2) presents different features. Let us explain our notations and then state our main results.

*Notation 1.4.* Let  $x$  be an element in the  $E_2$ -page of a certain spectral sequence. We let  $[x]_k$  denote the homology class of  $x$  on the  $E_k$ -page, where  $[x]_2$  is understood to be  $x$ . We say  $x$  can be lifted to the  $E_r$ -page if  $[x]_r$  is well defined, i.e., if  $d_i([x]_i) = 0$  for each  $2 \leq i < r$ .

**Theorem 1.5.** *Let  $p$  be an odd prime. Let  $z \in Ext_{\mathcal{A}_*}^{s+k,t+k}(\mathbb{F}_p, \mathbb{F}_p)$  be an element in the  $E_2$ -page of the ASS which is detected by  $x \in Ext_{P_*}^{s,t}(\mathbb{F}_p, I^k/I^{k+1})$  with  $s < 2p - 2$ . Let  $r$  be an integer such that  $r + k \leq 2p - 2$  and  $z$  can be lifted to the  $E_r$ -page. Then we can write  $d_r^{Adams}([z]_r) = [w]_r$ ,  $d_r^{alg}([x]_r) = [y]_r$ , where  $w \in Ext_{\mathcal{A}_*}^{s+k+r,t+k+r-1}(\mathbb{F}_p, \mathbb{F}_p)$  can be detected by  $y \in Ext_{P_*}^{s+1,t}(\mathbb{F}_p, I^{k+r-1}/I^{k+r})$ .*

*Remark 1.6.* In the motivic context, we say  $z \in Ext_{\mathcal{A}_*}(\mathbb{F}_p, \mathbb{F}_p)$  is detected by  $x \in Ext_{P_*}(\mathbb{F}_p, I^k/I^{k+1})$  if and only if  $z$  lifts to  $\tilde{z} \in Ext_{\mathcal{A}_{*,*}}(\mathbb{F}_p[\tau], \mathbb{F}_p[\tau])$  which maps to  $x \in Ext_{\mathcal{A}_{*,*}}(\mathbb{F}_p[\tau], \mathbb{F}_p) \cong Ext_{P_*}(\mathbb{F}_p, I^k/I^{k+1})$ , as shown in diagram (1.2).

Theorem 1.5 shows, in a certain range, the higher Adams differentials agree with their corresponding higher algebraic Novikov differentials. It is worth pointing out that the result could fail for  $r \geq 2p - 1 - k$ . For example, let  $p \geq 5$ . The  $k$ -value for  $b_n \in Ext_{\mathcal{A}_*}^{*,*}(\mathbb{F}_p, \mathbb{F}_p)$  is 0. It is proved that  $b_n$  has nontrivial Adams differential [14]

$$(1.3) \quad d_{2p-1}^{Adams}(b_n) = h_0 b_{n-1}^p.$$

However, the corresponding algebraic Novikov differential is trivial

$$(1.4) \quad d_{2p-1}^{alg}(b_n) = 0.$$

We also have the following result comparing the length of nontrivial algebraic Novikov differentials and Adams differentials.

**Theorem 1.7.** *Let  $p$  be an odd prime. Let  $z \in Ext_{\mathcal{A}_*}^{s+k,t+k}(\mathbb{F}_p, \mathbb{F}_p)$  be an element detected by  $x \in Ext_{P_*}^{s,t}(\mathbb{F}_p, I^k/I^{k+1})$  with  $s < 2p - 2$ . If  $x$  is not a permanent cycle in the algNSS, then  $z$  is not a permanent cycle in the ASS. Moreover, let  $r$  (resp.  $r'$ ) be the largest number  $n$  such that  $x$  (resp.  $z$ ) can be lifted to the  $E_n$ -page, then  $r \geq r'$ .*

**Organization of the paper.** In Section 2, we review some basic results in motivic homotopy theory as well as the “cofiber of  $\tau$ ” method of Gheorghie, Isaksen, Wang, and Xu to compare higher algebraic Novikov differentials with higher Adams differentials. In Section 3, we give the proofs for Theorem 1.5 and Theorem 1.7.

**Acknowledgments.** We would like to thank the anonymous referee for their constructive suggestions and detailed comments in revising this paper. The authors are supported by the National Natural Science Foundation of China (No. 12271183). The second named author is also supported by the National Natural Science Foundation of China (No. 12001474; 12261091).

## 2. COMPARING $d_r^{Adams}$ WITH $d_r^{alg}$

Motivic homotopy theory originated out of the work of Voevodsky and Morel [12, 16, 17]. Motivic homotopy theory can be viewed as a successful application of abstract homotopy theory to algebraic geometry and number theory. From categorical and computational perspectives, the motivic stable homotopy category behaves very similarly to the classical stable homotopy category. For example, there are motivic spheres, motivic homotopy and homology groups, motivic Eilenberg-MacLane spectra, and motivic Steenrod algebras analogous to the classical ones. For readers not familiar with motivic homotopy theory, [4, 11, 18] are some helpful references.

In this paper, we choose to work over the field  $\mathbb{C}$  of complex numbers and assume  $p$  is an odd prime. Under this setting, we have explicit formulas for the mod  $p$  motivic Eilenberg-MacLane spectra  $H\mathbb{F}_p^{mot}$  and motivic dual Steenrod algebra  $\mathcal{A}_{*,*}^{\mathbb{C}}$ .

**Proposition 2.1** ([16, 17]). *Let  $p$  be an odd prime. We have*

$$H\mathbb{F}_p^{mot}_{*,*} = \mathbb{F}_p[\tau],$$

where  $\tau$  has bi-degree  $(0, -1)$ , and that

$$\mathcal{A}_{*,*}^{\mathbb{C}} = \mathbb{F}_p[\tau] \otimes \mathbb{F}_p[t_1, t_2, \dots] \otimes E[\tau_0, \tau_1, \dots],$$

where  $\tau$  has bi-degree  $(0, -1)$ ,  $t_i$  has bi-degree  $(2(p^i - 1), p^i - 1)$ , and  $\tau_i$  has bi-degree  $(2(p^i - 1) + 1, p^i - 1)$ .

Under this setting,  $\mathcal{A}_{*,*}^{\mathbb{C}} \cong \mathbb{F}_p[\tau] \otimes_{\mathbb{F}_p} \mathcal{A}_*$  is just the classical dual Steenrod algebra tensoring the new coefficient.

There is a motivic analog of the classical Adams spectral sequence called the motivic Adams spectral sequence (mASS) (see [4, 6]).

**Proposition 2.2.** *There is a motivic Adams spectral sequence which converges to the bi-graded homotopy groups of the  $H\mathbb{F}_p^{mot}$ -completed motivic sphere  $\widehat{S}^{0,0}$ . The mASS has  $E_2$ -page*

$$E_2^{s,t,u}(S) = Ext_{\mathcal{A}_{*,*}^{\mathbb{C}}}^{s,t,u}(\mathbb{F}_p[\tau], \mathbb{F}_p[\tau]) \implies \pi_{t-s,u}(\widehat{S}^{0,0}),$$

and differentials

$$d_r : E_r^{s,t,u}(S) \rightarrow E_r^{s+r,t+r-1,u}(S).$$

For  $x \in E_r^{s,t,u}(S)$ , we call  $s$  its homological degree,  $t$  the inner degree, and  $u$  the motivic weight.

There is a topological realization functor  $Re : SH^{\text{mot}} \rightarrow SH$  (see [3, 12]) from the motivic stable homotopy category  $SH^{\text{mot}}$  to the classical stable homotopy category  $SH$ . This functor maps the motivic sphere  $S^{a,b}$  to classical sphere  $S^a$  and maps the motivic Eilenberg-MacLane spectrum  $H\mathbb{F}_p^{\text{mot}}$  to the classical Eilenberg-MacLane spectrum  $H\mathbb{F}_p$ . For the other direction, there is also a constant embedding functor  $C : SH \rightarrow SH^{\text{mot}}$ . We have  $Re \circ C = id$ .

The functor  $Re$  induces a map  $\phi$  from the mASS of the motivic sphere to the classical ASS of the classical sphere.

$$(2.1) \quad \begin{array}{ccc} E_r^{s,t,u}(S) & \xrightarrow{\phi_r} & E_r^{s,t}(S) \\ d_r^{mAdams} \downarrow & & \downarrow d_r^{Adams} \\ E_r^{s+r,t+r-1,u}(S) & \xrightarrow{\phi_r} & E_r^{s+r,t+r-1}(S) \end{array}$$

The effect of  $\phi_r$  is inverting  $\tau$  (see [3, 4, 7]).

**Proposition 2.3** ([4, 7]). *Let  $p$  be an odd prime. There is an isomorphism*

$$Ext_{\mathcal{A}_{*,*}^{\mathbb{C}}}^{s,t,*}(\mathbb{F}_p[\tau], \mathbb{F}_p[\tau]) \cong \mathbb{F}_p[\tau] \otimes_{\mathbb{F}_p} Ext_{\mathcal{A}_*}^{s,t}(\mathbb{F}_p, \mathbb{F}_p).$$

Moreover, after inverting  $\tau$ , the mASS of the motivic sphere becomes isomorphic to the classical ASS tensored over  $\mathbb{F}_p$  with  $\mathbb{F}_p[\tau, \tau^{-1}]$ .

The element  $\tau$  can be lifted to a map  $\tau : \widehat{S^{0,-1}} \rightarrow \widehat{S^{0,0}}$  between  $H\mathbb{F}_p^{\text{mot}}$ -completed motivic spectra. We denote the associated cofiber sequence as

$$\widehat{S^{0,-1}} \xrightarrow{\tau} \widehat{S^{0,0}} \xrightarrow{i} \widehat{S^{0,0}}/\tau.$$

The mASS for  $\widehat{S^{0,0}}/\tau$  has  $E_2$ -page  $E_2^{s,t,u}(C\tau) = Ext_{\mathcal{A}_{*,*}^{\mathbb{C}}}^{s,t,u}(\mathbb{F}_p[\tau], \mathbb{F}_p)$ . To avoid potential confusions with the differentials in the mASS for  $\widehat{S^{0,0}}$ , we denote the differentials in the mASS for  $\widehat{S^{0,0}}/\tau$  as  $\delta_r^{mAdams}$ . The map  $i$  induces a map  $\psi$  from the mASS for  $\widehat{S^{0,0}}$  to the mASS for  $\widehat{S^{0,0}}/\tau$ .

$$(2.2) \quad \begin{array}{ccc} E_r^{s,t,u}(C\tau) & \xleftarrow{\psi_r} & E_r^{s,t,u}(S) \\ \delta_r^{mAdams} \downarrow & & \downarrow d_r^{mAdams} \\ E_r^{s+r,t+r-1,u}(C\tau) & \xleftarrow{\psi_r} & E_r^{s+r,t+r-1,u}(S) \end{array}$$

The effect of  $\psi$  is just sending  $\tau$  to 0 (see [5, 8]).

Finally, Gheorghe-Wang-Xu [5] proved there is an isomorphism  $\kappa$  between the mASS for  $\widehat{S^{0,0}}/\tau$  and the regraded algebraic Novikov spectral sequence.

$$(2.3) \quad \begin{array}{ccc} \bar{E}_r^{s+2u-t, 2u, t-2u} & \xrightarrow[\cong]{\kappa_r} & E_r^{s,t,u}(C\tau) \\ d_r^{alg} \downarrow & & \downarrow \delta_r^{Adams} \\ \bar{E}_r^{s+2u-t+1, 2u, t-2u+r-1} & \xrightarrow[\cong]{\kappa_r} & E_r^{s+r,t+r-1,u}(C\tau) \end{array}$$

We can summarize these three comparison maps in the following diagram.

(2.4)

$$\begin{array}{ccccccc}
\bar{E}_r^{s+2u-t, 2u, t-2u} & \xrightarrow[\cong]{\kappa_r} & E_r^{s, t, u}(C\mathcal{T}) & \xleftarrow{\psi_r} & E_r^{s, t, u}(S) & \xrightarrow{\phi_r} & E_r^{s, t}(S) \\
d_r^{alg} \downarrow & & \delta_r^{m\text{ Adams}} \downarrow & & d_r^{m\text{ Adams}} \downarrow & & d_r^{Adams} \downarrow \\
\bar{E}_r^{s+2u-t+1, 2u, t-2u+r-1} & \xrightarrow[\cong]{\kappa_r} & E_r^{s+r, t+r-1, u}(C\mathcal{T}) & \xleftarrow{\psi_r} & E_r^{s+r, t+r-1, u}(S) & \xrightarrow{\phi_r} & E_r^{s+r, t+r-1}(S)
\end{array}$$

The diagram (2.4) provides a zig-zag way to compare higher Adams differentials  $d_r^{Adams}$  with their corresponding higher algebraic Novikov differentials  $d_r^{alg}$ .

### 3. PROOF OF THEOREMS 1.5, 1.7

We discuss several lemmas before we prove our main results.

**Lemma 3.1.** *Let  $p$  be an odd prime, denote  $q = 2(p-1)$ . Given  $s, t \geq 0$ , we denote  $C_{s,t} = \{i \in \mathbb{Z} \mid i \equiv t \pmod{q}, 0 \leq i \leq s, t\}$ . Then we have the following direct sum decomposition of the classical Adams  $E_2$ -terms.*

$$Ext_{A_*}^{s,t}(\mathbb{F}_p, \mathbb{F}_p) \cong \bigoplus_{i \in C_{s,t}} Ext_{P_*}^{s-i, t-i}(\mathbb{F}_p, I^i/I^{i+1})$$

*Proof.* For odd prime  $p$ , the Cartan-Eilenberg spectral sequence collapses from  $E_2$ -page with no nontrivial extensions [15, Theorem 4.4.3]. Hence we have

$$Ext_{A_*}^{s,t}(\mathbb{F}_p, \mathbb{F}_p) \cong \bigoplus_{i \in \mathbb{Z}} Ext_{P_*}^{s-i, t-i}(\mathbb{F}_p, I^i/I^{i+1}).$$

Note the inner degrees  $|v_n| = |t_n| = 2(p^n - 1)$  are all multiples of  $q$ . In order for  $Ext_{P_*}^{s-i, t-i}(\mathbb{F}_p, I^i/I^{i+1})$  to be nontrivial, we need  $i, s-i, t-i \geq 0$ , and that  $t-i \equiv 0 \pmod{q}$ . Hence  $i$  needs to be in the set  $C_{s,t}$ .  $\square$

*Remark 3.2.* It is worth pointing out that the Adams differential  $d_2^{Adams}$  may not respect this decomposition.

*Notation 3.3.* Let  $z \in Ext_{A_*}^{s,t}(\mathbb{F}_p, \mathbb{F}_p)$  be an element in the Adams  $E_2$ -page. We let  $\tilde{z}$  denote the element  $1 \otimes z \in \mathbb{F}_p[\tau] \otimes_{\mathbb{F}_p} Ext_{A_*}^{s,t}(\mathbb{F}_p, \mathbb{F}_p) \cong Ext_{A_*^c}^{s,t,*}(\mathbb{F}_p[\tau], \mathbb{F}_p[\tau])$  in the  $E_2$ -page of the mASS of the motivic sphere.

Note we have  $\phi_2(\tilde{z}) = z$ . If  $z$  and  $\tilde{z}$  can both be lifted to the  $E_r$ -pages of the respected spectral sequences for some  $r$ . Then  $\phi_r([\tilde{z}]_r) = [z]_r$ .

**Lemma 3.4.** *Let  $z$  be an element in  $Ext_{A_*}^{s,t}(\mathbb{F}_p, \mathbb{F}_p)$  which is detected by  $x \in Ext_{P_*}^{s-k, t-k}(\mathbb{F}_p, I^k/I^{k+1})$ . Then  $\tilde{z} \in Ext_{A_*^c}^{s,t,*}(\mathbb{F}_p[\tau], \mathbb{F}_p[\tau])$  has motivic weight  $\frac{t-k}{2}$ .*

*Proof.* For notation simplicity, we let  $t(-)$  denote the inner degree and let  $u(-)$  denote the motivic weight of an element in  $Ext_{A_*^c}^{*,*,*}(\mathbb{F}_p[\tau], \mathbb{F}_p[\tau])$ . By Proposition 2.1, we have  $u(\tilde{t}_i) = \frac{1}{2}t(\tilde{t}_i)$  for  $i \geq 1$ , and  $u(\tilde{\tau}_i) = \frac{1}{2}t(\tilde{\tau}_i) - \frac{1}{2}$  for  $i \geq 0$ .

Since  $z$  is detected by  $x \in Ext_{P_*}^{s-k, t-k}(\mathbb{F}_p, I^k/I^{k+1})$ , the number of  $\tau_i$ 's in the expression of  $z$  is just  $k$ . Hence we conclude  $u(\tilde{z}) = \frac{1}{2}t(\tilde{z}) - \frac{1}{2}k = \frac{t-k}{2}$ .  $\square$

**Lemma 3.5.** *Let  $z \in Ext_{\mathcal{A}_*}^{s+k, t+k}(\mathbb{F}_p, \mathbb{F}_p)$  be an element which is detected by  $x \in Ext_{\mathcal{P}_*}^{s, t}(\mathbb{F}_p, I^k/I^{k+1})$ . (i) If  $\tilde{z}$  can be lifted to the  $E_r$ -page for some  $r \geq 2$ , then  $z$  can also be lifted to the  $E_r$ -page. (ii) Further assume  $s < 2p - 2$ . Then  $z$  can be lifted to the  $E_r$ -page implies  $\tilde{z}$  can also be lifted to the  $E_r$ -page.*

*Proof.* (i) By assumption,  $d_i^{mAdams}([\tilde{z}]_i) = 0$  for each  $2 \leq i < r$ . One can inductively show that  $d_i^{mAdams}([z]_i) = 0$  (hence  $[z]_{i+1}$  is well defined) for each  $2 \leq i < r$  by commutativity of diagram (2.1).

(ii) Assume, for the sake of contradiction, that there exists  $2 \leq r_1 < r$  such that  $d_{r_1}^{mAdams}([\tilde{z}]_{r_1}) \neq 0$ . We claim that there exists a nontrivial differential  $d_j^{mAdams}([u]_j) \neq 0$ , where  $u \in Ext_{\mathcal{A}_{*,*}^c}^{s+b, t+b, \frac{t}{2}-a}(\mathbb{F}_p[\tau], \mathbb{F}_p[\tau])$  is non- $\tau$ -divisible,  $2 \leq j < r_1$ , and  $a > 0$ .

We let  $[v_1]_{r_1}$  denote  $d_{r_1}^{mAdams}([\tilde{z}]_{r_1}) \neq 0$ . By the commutativity of diagram (2.1), we have  $\phi_{r_1}([v_1]_{r_1}) = d_{r_1}^{mAdams}([z]_{r_1}) = 0$ . So  $[v_1]_{r_1}$  is  $\tau$ -torsion. Let  $a_1 > 0$  be the smallest integer such that  $\tau^{a_1}[v_1]_{r_1} = 0$ . Then there exists a nontrivial differential  $d_{r_2}^{mAdams}([u_1]_{r_2}) = \tau^{a_1}[v_1]_{r_2} \neq 0$  with  $2 \leq r_2 < r_1$ . The element  $[u_1]_{r_2}$  is not divisible by  $\tau$  on the  $E_{r_2}$ -page. Otherwise, we have  $d_{r_2}^{mAdams}([u_1]_{r_2}/\tau) = \tau^{a_1-1}[v_1]_{r_2}$ , which contradicts the definition of  $a_1$ .

By Lemma 3.4, we have  $[\tilde{z}]_{r_1} \in E_{r_1}^{s+k, t+k, \frac{t}{2}}(S)$ . This implies

$$[v_1]_{r_1} = d_{r_1}^{mAdams}([\tilde{z}]_{r_1}) \in E_{r_1}^{s+k+r_1, t+k+r_1-1, \frac{t}{2}}(S).$$

Comparing the degrees, we have

$$\tau^{a_1}[v_1]_{r_2} \in E_{r_2}^{s+k+r_1, t+k+r_1-1, \frac{t}{2}-a_1}(S), [u_1]_{r_2} \in E_{r_2}^{s+k+r_1-r_2, t+k+r_1-r_2, \frac{t}{2}-a_1}(S).$$

If  $u_1$  is non- $\tau$ -divisible, we can take  $d_{r_2}^{mAdams}([u_1]_{r_2}) \neq 0$  as the claimed differential.

Otherwise, since  $[u_1]_{r_2}$  is not divisible by  $\tau$  on the  $E_{r_2}$ -page, we conclude  $u_1/\tau$  does not lift to the  $E_{r_2}$ -page. Then there exists differential  $d_{r_3}^{mAdams}([u_1/\tau]_{r_3}) = [v_2]_{r_3} \neq 0$  with  $2 \leq r_3 < r_2$ .

By the commutativity of diagram (2.1), we have

$$\phi_{r_3}([v_2]_{r_3}) = d_{r_3}^{mAdams}(\phi_{r_3}([u_1/\tau]_{r_3})) = d_{r_3}^{mAdams}(\phi_{r_3}([u_1]_{r_3})) = \phi_{r_3}(d_{r_3}^{mAdams}([u_1]_{r_3})),$$

and  $d_{r_3}^{mAdams}([u_1]_{r_3}) = 0$  since  $u_1$  can be lifted to the  $E_{r_2}$ -page. So  $[v_2]_{r_3}$  is  $\tau$ -torsion. Let  $a_2 > 0$  be the smallest integer such that  $\tau^{a_2}[v_2]_{r_3} = 0$ . Then there exists a nontrivial differential  $d_{r_4}^{mAdams}([u_2]_{r_4}) = \tau^{a_2}[v_2]_{r_4} \neq 0$  with  $2 \leq r_4 < r_3$ . The element  $[u_2]_{r_4}$  is not divisible by  $\tau$  on the  $E_{r_4}$ -page.

For the degrees, we have

$$\begin{aligned} [u_1/\tau]_{r_3} &\in E_{r_3}^{s+k+r_1-r_2, t+k+r_1-r_2, \frac{t}{2}-(a_1-1)}(S), \\ [v_2]_{r_3} &\in E_{r_3}^{s+k+r_1-r_2+r_3, t+k+r_1-r_2+r_3-1, \frac{t}{2}-(a_1-1)}(S), \\ \tau^{a_2}[v_2]_{r_4} &\in E_{r_4}^{s+k+r_1-r_2+r_3, t+k+r_1-r_2+r_3-1, \frac{t}{2}-(a_1-1)-a_2}(S), \\ [u_2]_{r_4} &\in E_{r_4}^{s+k+r_1-r_2+r_3-r_4, t+k+r_1-r_2+r_3-r_4, \frac{t}{2}-(a_1-1)-a_2}(S). \end{aligned}$$

If  $u_2$  is non- $\tau$ -divisible, we can take  $d_{r_4}^{mAdams}([u_2]_{r_4}) \neq 0$  as the claimed differential. Otherwise, we can repeat the process and obtain  $u_3, u_4, \dots$ . Note there are only finitely many integers between 2 and  $r_1$ . After repeating this process several times,

we will eventually obtain a desired nontrivial differential  $d_j^{mAdams}([u]_j) \neq 0$ , where  $u \in Ext_{\mathcal{A}_{*,*}^{\mathbb{C}}}^{s+b, t+b, \frac{t}{2}-a}(\mathbb{F}_p[\tau], \mathbb{F}_p[\tau])$  is non- $\tau$ -divisible,  $2 \leq j < r_1$ , and  $a > 0$ .

By Proposition 2.3 and Lemma 3.1, we can write

$$u = \sum_{i \in C_{s+b, t+b}} \tau^{n_i} \tilde{u}_i,$$

where  $u_i$  can be detected by some  $m_i \in Ext_{P_*}^{s+b-i, t+b-i}(\mathbb{F}_p, I^i/I^{i+1})$ . Comparing the motivic weights using Lemma 3.4, we get  $n_i = \frac{b-i+2a}{2} \geq 0$  for  $\tilde{u}_i \neq 0$ .

Note  $u$  is non- $\tau$ -divisible. This implies the  $n_i = 0$  term is nontrivial. In particular, we have  $b+2a \in C_{s+b, t+b}$ . This implies

$$(3.1) \quad b+2a \leq s+b, \quad b+2a \equiv t+b \pmod{q}.$$

Note  $x \in Ext_{P_*}^{s,t}(\mathbb{F}_p, I^k/I^{k+1})$  implies  $t \equiv 0 \pmod{q}$ . In summary, we have

$$(3.2) \quad 0 < 2a \leq s, \quad 2a \equiv 0 \pmod{q}.$$

Then  $s \geq 2a \geq 2(p-1)$ , this contradicts  $s < 2p-2$ . Thus we have proved (ii).  $\square$

Now we proceed to prove Theorem 1.5 and Theorem 1.7.

*Proof of Theorem 1.5.* Our strategy is to compare the differentials via diagram (2.4). As we will see, with the given assumptions, diagram (2.4) can be specialized to the following diagram.

$$(3.3) \quad \begin{array}{ccccccc} [x]_r & \xrightarrow{\kappa_r} & [\tilde{z}]_r & \xleftarrow{\psi_r} & [\tilde{z}]_r & \xrightarrow{\phi_r} & [z]_r \\ d_r^{alg} \downarrow & & \downarrow \delta_r^{mAdams} & & \downarrow d_r^{mAdams} & & \downarrow d_r^{Adams} \\ [y]_r & \xrightarrow{\kappa_r} & [\tilde{w}]_r & \xleftarrow{\psi_r} & [\tilde{w}]_r & \xrightarrow{\phi_r} & [w]_r \end{array}$$

By Lemma 3.5,  $[\tilde{z}]_r$  is well defined and we have  $\phi_r([\tilde{z}]_r) = [z]_r$ . By Lemma 3.4, we have  $[\tilde{z}]_r \in E_r^{s+k, t+k, \frac{t}{2}}(S)$ . Then we can write  $d_r^{mAdams}([\tilde{z}]_r) = [u]_r$ , where  $u \in Ext_{\mathcal{A}_{*,*}^{\mathbb{C}}}^{s+k+r, t+k+r-1, \frac{t}{2}}(\mathbb{F}_p[\tau], \mathbb{F}_p[\tau])$ . By Proposition 2.3 and Lemma 3.1, we can write

$$(3.4) \quad u = \sum_{i \in C_{s+k+r, t+k+r-1}} \tau^{n_i} \tilde{u}_i,$$

where  $u_i$  can be detected by some  $m_i \in Ext_{P_*}^{s+k+r-i, t+k+r-1-i}(\mathbb{F}_p, I^i/I^{i+1})$ . Comparing the motivic weights using Lemma 3.4, we get  $n_i = \frac{k+r-1-i}{2}$  for  $\tilde{u}_i \neq 0$ .

Since  $n_i \geq 0$ , this forces  $i \leq k+r-1$ . By definition,  $i \in C_{s+k+r, t+k+r-1}$  implies

$$(3.5) \quad i \geq 0, \quad i \equiv t+k+r-1 \pmod{q},$$

where we denote  $q = 2(p-1)$ . Moreover, since  $x \in Ext_{P_*}^{s,t}(\mathbb{F}_p, I^k/I^{k+1})$ , we have  $t \equiv 0 \pmod{q}$ . In summary, we have

$$(3.6) \quad 0 \leq i \leq k+r-1, \quad i \equiv k+r-1 \pmod{q}$$

for nontrivial  $\tilde{u}_i$ .

By assumption, we have  $0 < k+r-1 < q$ . Then (3.6) forces  $i = k+r-1$ , the corresponding  $n_i = 0$ . So we can rewrite (3.4) as  $u = \tilde{w}$ , where we let  $w$  denote  $u_{k+r-1}$ . We also let  $y$  denote  $m_{k+r-1}$  detecting  $w$ .



We have  $d_r^{mAdams}([\tilde{z}]_r) = [\tilde{w}]_r$ . Note  $\phi_r([\tilde{w}]_r) = [w]_r$ . The commutativity of diagram (2.1) implies  $d_r^{Adams}([z]_r) = [w]_r$ .

Note  $[\tilde{z}]_r$  and  $[\tilde{w}]_r$  are non- $\tau$ -divisible,  $\psi$  sends  $[\tilde{z}]_r$  and  $[\tilde{w}]_r$  to the corresponding elements in  $E_r^{*,*,*}(C\tau)$  of the same form, which we abuse the notation and still denote by  $[\tilde{z}]_r$  and  $[\tilde{w}]_r$  respectively. The commutativity of diagram (2.2) implies  $\delta_r^{mAdams}([\tilde{z}]_r) = [\tilde{w}]_r$ .

Finally, the isomorphism  $\kappa$  associates  $\tilde{z}$  with  $x$  and  $\tilde{w}$  with  $y$ . Hence  $\kappa_r([x]_r) = [\tilde{z}]_r$ ,  $\kappa_r([y]_r) = [\tilde{w}]_r$ , and  $d_r^{alg}([x]_r) = [y]_r$ .

Now we have completed diagram (3.3). The results of the theorem follow directly.  $\square$

*Proof of Theorem 1.7.* We can prove the following equivalent statement: suppose  $r \geq 2$  is an integer such that  $z$  can be lifted to the  $E_r$ -page (of the ASS). Then  $x$  can also be lifted to the  $E_r$ -page (of the algNSS).

We will establish the statement using an inductive approach. By definition,  $x$  can be lifted to the  $E_2$ -page. Next, assuming that  $x$  can be lifted to the  $E_i$ -page with  $2 \leq i < r$ , we need to show that  $d_i^{alg}([x]_i) = 0$ . This will imply that  $x$  can be lifted to the  $E_{i+1}$ -page.

We study the differential  $d_i^{alg}([x]_i)$  via diagram (2.4). As we will see, with the given assumptions, diagram (2.4) can be specialized to the following diagram.

$$(3.7) \quad \begin{array}{ccccccc} [x]_i & \xrightarrow{\kappa_i} & [\tilde{z}]_i & \xleftarrow{\psi_i} & [\tilde{z}]_i & \xrightarrow{\phi_i} & [z]_i \\ d_i^{alg} \downarrow & & \downarrow \delta_i^{mAdams} & & \downarrow d_i^{mAdams} & & \downarrow d_i^{Adams} \\ 0 & \xrightarrow{\kappa_i} & 0 & \xleftarrow{\psi_i} & 0 & \xrightarrow{\phi_i} & 0 \end{array}$$

By Lemma 3.5,  $\tilde{z}$  can be lifted to the  $E_r$ -page of the mASS of the sphere. Hence  $d_i^{mAdams}([\tilde{z}]_i) = 0$ . Note  $[\tilde{z}]_i$  is non- $\tau$ -divisible, so  $\psi$  sends  $[\tilde{z}]_i$  to the corresponding element in  $E_r^{*,*,*}(C\tau)$  of the same form, which we abuse the notation and still denote by  $[\tilde{z}]_i$ . The commutativity of diagram (2.2) implies  $\delta_i^{mAdams}([\tilde{z}]_i) = 0$ . Finally, using the isomorphism  $\kappa$ , we deduce  $d_i^{alg}([x]_i) = 0$ .  $\square$

## REFERENCES

- [1] Michael Andrews and Haynes Miller. Inverting the Hopf map. *J. Topol.*, 10(4):1145–1168, 2017.
- [2] Henri Cartan and Samuel Eilenberg. *Homological algebra*. Princeton University Press, Princeton, N. J., 1956.
- [3] Daniel Dugger and Daniel C. Isaksen. Topological hypercovers and  $\mathbb{A}^1$ -realizations. *Math. Z.*, 246(4):667–689, 2004.
- [4] Daniel Dugger and Daniel C. Isaksen. The motivic Adams spectral sequence. *Geom. Topol.*, 14(2):967–1014, 2010.
- [5] Bogdan Gheorghe, Guozhen Wang, and Zhouli Xu. The special fiber of the motivic deformation of the stable homotopy category is algebraic. *Acta Math.*, 226(2):319–407, 2021.
- [6] P. Hu, I. Kriz, and K. Ormsby. Convergence of the motivic Adams spectral sequence. *J. K-Theory*, 7(3):573–596, 2011.
- [7] Daniel C. Isaksen. Stable stems. *Mem. Amer. Math. Soc.*, 262(1269):viii+159, 2019.
- [8] Daniel C Isaksen, Guozhen Wang, and Zhouli Xu. Stable homotopy groups of spheres: From dimension 0 to 90. *Publications mathématiques de l’IHÉS*, 137(1):107–243, 2023.
- [9] Haynes R. Miller. *Some algebraic aspects of the Adams–Novikov spectral sequence*. PhD thesis, Princeton University, 1974.

- [10] Haynes R. Miller. On relations between Adams spectral sequences, with an application to the stable homotopy of a Moore space. *J. Pure Appl. Algebra*, 20(3):287–312, 1981.
- [11] Fabien Morel. An introduction to  $\mathbb{A}^1$ -homotopy theory. In *Contemporary developments in algebraic K-theory*, ICTP Lect. Notes, XV, pages 357–441. Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2004.
- [12] Fabien Morel and Vladimir Voevodsky.  $\mathbb{A}^1$ -homotopy theory of schemes. *Inst. Hautes Études Sci. Publ. Math.*, (90):45–143 (2001), 1999.
- [13] S. P. Novikov. Methods of algebraic topology from the point of view of cobordism theory. *Izv. Akad. Nauk SSSR Ser. Mat.*, 31:855–951, 1967.
- [14] Douglas C. Ravenel. The non-existence of odd primary Arf invariant elements in stable homotopy. *Math. Proc. Cambridge Philos. Soc.*, 83(3):429–443, 1978.
- [15] Douglas C. Ravenel. *Complex cobordism and stable homotopy groups of spheres*, volume 121 of *Pure and Applied Mathematics*. Academic Press, Inc., Orlando, FL, 1986.
- [16] Vladimir Voevodsky. Reduced power operations in motivic cohomology. *Publ. Math. Inst. Hautes Études Sci.*, (98):1–57, 2003.
- [17] Vladimir Voevodsky. Motivic Eilenberg-MacLane spaces. *Publ. Math. Inst. Hautes Études Sci.*, (112):1–99, 2010.
- [18] Vladimir Voevodsky, Oliver Röndigs, and Paul Arne Østvær. Voevodsky’s Nordfjordeid lectures: motivic homotopy theory. In *Motivic homotopy theory*, Universitext, pages 147–221. Springer, Berlin, 2007.
- [19] Xiangjun Wang, Yaxing Wang, and Yu Zhang. Some nontrivial secondary Adams differentials on the fourth line. *New York J. Math.*, 29:687–707, 2023.

DEPARTMENT OF MATHEMATICS, NANKAI UNIVERSITY, NO.94 WEIJIN ROAD, TIANJIN 300071, PEOPLE’S REPUBLIC OF CHINA

*Email address:* `xjwang@nankai.edu.cn`

CENTER FOR APPLIED MATHEMATICS, TIANJIN UNIVERSITY, NO.92 WEIJIN ROAD, TIANJIN 300072, PEOPLE’S REPUBLIC OF CHINA

*Email address:* `zhang.4841@buckeyemail.osu.edu`